



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

the Eocene of Africa: "Systematisch betrachtet trenne ich die Archaeoceti vollständig von den Cetaceen und schliesse sie als Untergruppe an die Creodontier an, während die übrigen Walthiere nach wie vor eine selbständige Gruppe bilden, die so lange beliebig in der Systematik eingeschaltet werden kann, bis wir deren Stammesgeschichte kennen;" and his conclusion that these animals belong among the early carnivora seems well substantiated by him.

---

S. W. W.

"*Teleorhinus browni*—a New Teleosaur in the Fort Benton,"  
*Bulletin of the American Museum of Natural History*, Vol. XX  
(1904), p. 239.

Another discovery of much interest is that of a true teleosaur crocodile from the Benton Cretaceous, by Professor Osborn. That the teleosaurs should occur in America is perhaps not so remarkable as their occurrence in the Upper Cretaceous, all the forms hitherto known being from the Jurassic and Wealden of Europe. The specimen upon which Professor Osborn bases his new genus *Teleorhinus* has a skull one meter in length, and typical teleosaurian vetebrae, the latest biconcave crocodile vertebrae known.

S. W. W.